

PATIENT COUNSELLING INFORMATION

Diastase & Pepsin Liquid

Aristozyme[®] Liquid

Read the entire leaflet carefully before you start taking this medicine:

- If you have any further questions, ask your doctor.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet.

1. What is Aristozyme[®] Liquid and what is it used for?

Aristozyme[®] Liquid is a preparation, which contains digestive enzyme Diastase for effective digestion of carbohydrates & Pepsin for digestion of proteins. It is indicated in dyspepsia, epigastric distress and indigestion.

2. What you need to know before you take Aristozyme[®] Liquid?

Do not take Aristozyme[®] Liquid if you are allergic (hypersensitive) to Diastase, Pepsin or any other ingredients of this medicine and tell your doctor.

Other medicines and Aristozyme[®] Liquid

Tell your doctor if you are taking, or have recently taken, or might take any other medicines.

Pregnancy, breast-feeding and fertility

If you are pregnant, planning to become pregnant or are breast-feeding, speak to your doctor before taking this medicine.

3. How to take Aristozyme[®] Liquid ?

Always take Aristozyme[®] Liquid exactly as your doctor has told you. If you are not sure, check with your doctor. Take Aristozyme[®] Liquid every day for as long as your doctor prescribes it.

If you take more Aristozyme[®] Liquid than you should

It is important to keep to the dose as prescribed by your doctor. If you have taken more Aristozyme[®] Liquid than you should and if you are concerned about an overdose you should consult your doctor, if possible you should take the medicine pack with you.

If you forget to take the Aristozyme[®] Liquid

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, take it as soon as you remember it and then take the next dose at the right time.

If you stop taking Aristozyme[®] Liquid

If you stop taking Aristozyme[®] Liquid, your symptoms may return. If your doctor decides to stop this medication, your dose may be decreased gradually over a few days. If you have any further questions on the use of this medicine, ask your doctor.

4. What are the possible side effects of Aristozyme[®] Liquid?

Like all medicines, this medicine can cause side effects, although not everybody gets them. Side-effects with Aristozyme[®] Liquid mostly occur at the start of treatment and usually disappear within a few days with continued treatment. Occasional side-effects include nausea, constipation, skin rash, and pruritus.

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet.

5. How to store Aristozyme[®] Liquid?

Do not take this medicine after the expiry date shown on the pack. The expiry date refers to the last day of that month.

Store in a cool & dark place.

Keep out of reach of children.

Date of revision: June 15, 2021

® Registered Trademark

For further information please write to :



Marketed by:
ARISTO Pharmaceuticals Private Limited.
Regd. Office: 12, J.N. Heredia Marg,
Mumbai-400 001.